

Tidal Barrier and the Asymptotic Mass of Proto Gas-Giant Planets

Ian Dobbs-Dixon¹, Shu Lin Li^{1,2}, and D. N. C. Lin¹

¹*Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA*

²*Department of Astronomy, Peking University, Beijing, China*

ABSTRACT

The extrasolar planets found with radial velocity surveys have masses ranging from several Earth to several Jupiter masses. According to the conventional sequential accretion scenario, these planets acquired super-earth cores prior to the onset of efficient gas accretion with rates that rapidly increase with their masses. In weak-line T-Tauri disks, mass accretion onto protoplanetary cores may eventually be quenched by a global depletion of gas, as in the case of Uranus and Neptune. However, such a mechanism is unlikely to have stalled the growth of some known planetary systems which contain relatively low-mass and close-in planets along with more massive and longer period companions. Here, we suggest a potential solution for this conundrum. In general, supersonic infall of surrounding gas onto a protoplanet is only possible interior to both of its Bondi and Roche radii. At a critical mass, a protoplanet's Bondi and Roche radii are equal to the disk thickness. Above this mass, the protoplanets' tidal perturbation induces the formation of a gap. Although the disk gas may continue to diffuse into the gap, the azimuthal flux across the protoplanets' Roche lobe is quenched. Using two different schemes, we present the results of numerical simulations and analysis to show that the accretion rate increases rapidly with the ratio of the protoplanet's Roche to Bondi radii or equivalently to the disk thickness. In regions with low geometric aspect ratios, gas accretion is quenched with relatively low protoplanetary masses. This effect is important for determining the gas-giant planets' mass function, the distribution of their masses within multiple planet systems around solar type stars, and for suppressing the emergence of gas-giants around low mass stars. Finally, we find that the accretion rate into the protoplanets' Roche lobe declines gradually on a characteristic timescale of a few Myr. During this final stage, the protracted decline in the accretion flow across their Roche lobe and onto their protoplanetary disks may lead to the formation and retention of their regular satellites.

Subject headings: planet formation, accretion

1. Introduction

In the conventional planet formation scenario (Safronov & Zvjagina 1969; Pollack et al. 1996; Ida & Lin 2004) heavy elements first condense into grains which then grow into planetesimals. In a gaseous environment, the hydrodynamic and tidal interaction between the disk gas and these planetesimals damps their eccentricity (Adachi et al. 1976; Ward 1998). Nevertheless, these planetesimals can perturb each other’s orbit and induce growth through coagulation (Aarseth et al. 1993; Palmer et al. 1993). This growth process is stalled with the emergence of protoplanetary embryos that have acquired all the residual planetesimals within their feeding zone, which is 3-5 times that of their Roche radius (Kokubo & Ida 1998). In protostellar disks with similar mass and temperature distributions as those inferred for the primordial solar nebula, the embryos attain isolation masses that are typically a few times that of the Moon at 1AU (Lissauer 1987; Ida & Lin 2004).

In principle, the protoplanetary embryos can accrete gas from the disk when they have attained sufficient mass such that their surface escape speed is larger than the sound-speed in the disk. However, the gas which falls onto the embryos also converts gravitational energy into heat which is then transported to the surface of the nascent disk (Bodenheimer & Pollack 1986). Unless the cores have acquired more than a few earth masses ($M_{\oplus}$), the envelopes around them are too tenuous to allow efficient radiative or convective transport (Pollack et al. 1996). Consequently, a quasi hydrostatic equilibrium is established and gas sediments onto the cores on a protracted cooling timescale.

Once the protoplanet acquires sufficient mass to allow gas to be freely accreted, it enters a dynamical gas accretion phase. During this stage, the cores’ gravity dominates the flow out to the Bondi radius (R_B), where the local escape speed becomes comparable to the gas sound-speed in the disk. This radius increases linearly with the total mass of the protoplanet, which includes that of the core and the envelope. The protoplanets’ Roche radius (R_R) also increases with the mass of the protoplanet, though the mass dependence is much weaker. Although R_B is generally smaller than R_R for low-mass protoplanets, R_B overtakes R_R for sufficiently large protoplanetary masses.

As gas within R_R collapses toward the cores, it is replenished by the diffusion of gas from beyond R_R . The rate of matter diffusion is generally assumed to be induced by turbulence, though the origin of such turbulence in protostellar disks at a few AU’s remains an outstanding issue. Protoplanets also exert a tidal perturbation on the disk with a strength that

increases with their mass. Around relatively massive protoplanets, this tidal perturbation is sufficiently strong to induce the formation of a gap in the disk near the protoplanets’ orbits (Goldreich & Tremaine 1979, 1980) and to quench the accretion (Lin & Papaloizou 1979, 1985). In relatively cold and weakly-viscous disks, the asymptotic mass of the protoplanets near the ice line in the solar nebula is comparable to that Jupiter (Lin & Papaloizou 1993; Bryden et al. 1999).

However, there are numerical simulations of disk-planet interaction which indicate that gas continues to flow into the gap after its formation conditions have been satisfied (Lubow et al. 1999; Kley et al. 2001; D’Angelo et al. 2002). In these simulations, the magnitude of both implicit and numerical viscosity is sufficiently large that the intrinsic redistribution of angular momentum in the disk gas is comparable to the tidal torque exerted by the embedded protoplanet. Several simulations also take into account the effect of the turbulent flow (Armitage 1998; Nelson & Papaloizou 2003; Laughlin et al. 2004). All these models show that gas continues to flow to the vicinity of the protoplanet until its asymptotic mass becomes several times that of the Jupiter.

Although radial velocity surveys have discovered many planets with masses comparable to Saturn, the mass of Saturn is well below that found in the aforementioned simulations. While Saturn may have formed during the late stages of protostellar evolution when the disk mass was severely reduced (Guillot 2005), such a scenario cannot account for the relatively low mass of several close-in and resonant planets since it is generally accepted that their kinematic configuration requires extensive migration through active protostellar disks (Lin et al. 1996). In this paper, we consider another effect which may limit the asymptotic mass of protoplanets involving the tidal barrier. Due to its multi-dimensional nature, the complex flow pattern is ideally analyzed with numerical simulations.

The earliest disk simulations did not have sufficient resolution to resolve flow near the protoplanet (Miki 1982). A novel approach was introduced by Korycansky & Papaloizou (1996) to analyze inviscid flow in a steady-state. Under these conditions, the potential vorticity (hereafter vortensity) and the Bernoulli energy are conserved. In their simulations, they focused their attention on a local patch close to the protoplanet by adopting a set of boundary conditions which approximate the nearly co-orbiting flow at large azimuthal distance with a Keplerian velocity and smooth surface density. The advantage of this approach is that the high resolution allows the flow pattern to be accurately simulated (Balmforth & Korycansky 2001). In the absence of viscosity, however, the stream lines around the protoplanets are not connected to that beyond the Roche radius.

Adopting the same boundary condition, Tanigawa & Watanabe (2002) carried out a series of high resolution time-dependent nonlinear simulations which clearly demonstrate the

presence of shock dissipation near the accreting protoplanet and generation/dissipation of vortensity across the shock. Nevertheless, between the shock waves, quasi steady flows can be established with conserved vortensity (ϖ) and Bernoulli energy E_B . They also demonstrate the possibility that a quasi steady-state may not be attainable under some circumstances. The minor deficiency of both these simulations was the artificial boundary condition at large azimuth from the protoplanet. We demonstrate here that the modification of the global disk structure can significantly alter the accretion rate onto the embedded protoplanets.

There are many 2D global simulations of protoplanet-disk interaction (Papaloizou & Terquem 2006, see the recent review by). These simulations show that tidal torque from a sufficiently massive embedded protoplanet not only induces dissipative shock in the disk but also leads to gap formation (Lin & Papaloizou 1993; Artymowicz & Lubow 1994; Bryden et al. 1999; Kley 1999, cf.). There are also 3D simulations that show gas flow into the Roche lobe of the protoplanet (Bate et al. 2003; Klahr & Kley 2006), and some of these simulations have sufficient resolution to resolve the flow near the protoplanet (D’Angelo et al. 2002). The accretion rate determined from these simulations suggests that despite a significant reduction in Σ near the orbit of the protoplanet, the growth timescale for Jupiter-mass protoplanets is shorter than the global-disk gas depletion timescale ($\tau_{\text{dep}} \sim 3 - 10$ Myr) inferred from the observation of protoplanetary disks (Haisch et al. 2001; Hartmann 2005; Silverstone et al. 2006). If the gap region is continually replenished by diffusion into the gap, the asymptotic mass of the protoplanets would well exceed the observed upper limit of the planetary mass function (Marcy et al. 2005).

Here, we explore the possibility that even if the gap is replenished with tenuous gas, only a small fraction of that gas may be accreted onto protoplanets with Roche radius (R_R) greater than the disk thickness H . The model parameters in many previous simulations are chosen to approximate those inferred from the minimum mass nebula model in which the ratio between the disk thickness and radius (H/a) is assumed to be around 0.07. In these simulations, H determines the magnitude of the sound-speed and therefore the pressure gradient in the radial direction in terms of the surface density distribution Σ . With this assumption, R_B becomes comparable to R_R for a protoplanet with $M_p \sim M_J$.

One exception to this set of parameters was the consideration of a perturbation by a one-third Neptune-mass planet in very thin/cold disk with $H/a \sim 0.02 - 0.03$ (Bryden et al. 2000). In this case, R_B is several times R_R and the surface density of the disk diminishes well below that of an unperturbed disk. Although gas was not allowed to accrete onto the core of the protoplanet, the accretion timescales inferred from these simulations is substantially longer than those inferred from the idealized Bondi analysis, the viscous diffusion of the disk, and the τ_{dep} inferred from the observed SED evolution of disks in young clusters.

The disk thickness (H) is determined by heating from both the intrinsic viscous dissipation and surface irradiation from the central star. The commonly assumed aspect ratio ($H/a \sim 0.07$) at the present-day distance of Jupiter (5.2AU) is determined under the assumption that the disk is heated to an equilibrium temperature by the proto-sun with the present-day luminosity of the Sun. Self consistent models indicate that the H/a at the snow line can be significantly smaller than that assumed in the solar nebula model (Garaud & Lin 2006). Taking this possibility into account, we suggest that, in the limit of $R_R > H$, a tidal barrier imposed by a protoplanet limits the gas flow from most regions inside the gap into the Roche lobe. Much of the mass that diffuses into the gap accumulates into the horseshoe orbits in the co-orbital region, but is not able to accrete onto the planet.

In order to examine the dominant physical effects which regulate protoplanets’ asymptotic masses, we calculate the accretion rates and flow dynamics through two sets of simulations. Since we are primarily interested in the limit $R_R > H$, the flow pattern is well approximated by an 2D simulation (D’Angelo et al. 2002). In §2, we present an analysis to highlight the nature and importance of the tidal barrier. Due to the complexity of the 2D flow, we cannot obtain a full set of analytic solutions. For illustration purposes, we also adopt a grossly simplified 1D approximation for the protoplanet’s tidal potential and obtain solutions to highlight the effect of the tidal barrier in quenching the accretion flow onto it.

In order to illustrate the structure of the flow, we present in §3, a set of high-resolution 2D hydrodynamic simulations to demonstrate that the flow into the Roche lobe is indeed quenched for protoplanets with $R_R > H$ despite diffusion of gas into the gap region. Although the 2D simulation provides a more realistic approximation for the multi-dimensional geometry of the potential field and the non-spherically symmetric flow pattern of the background gas, numerical simulations of this kind can only be carried out with a limited resolution for several dynamical timescales. However, the disk environment, as well as the mass growth and envelope structure of the protoplanet, all evolve on much longer timescales, especially during the epoch when the envelope undergoes a transition from quasi hydrostatic contraction to dynamical accretion. The computation of the late stages of giant planet formation also requires following the governing equations over large dynamical range in space and time. While the envelope around the core adjusts to quasi hydrostatic equilibrium, the flow around the planet’s Hills radius adjusts to a steady state. The technical challenges for 2D simulations covering more than four orders of magnitude in physical scales over such large timescales makes them impractical for this purpose. In §3.3, we show that the dominant physical effects generated by the 2D simulations can be quantitatively captured by a simple-to-use 1D scheme, with which we verify that the gas accretion is significantly suppressed in the limit $R_R > H$. Utilizing this model, we show that the flow of gas into the protoplanets’ Roche lobe declines and the protoplanets acquire asymptotic masses on Myr timescale. Finally, we

summarize our results and discuss their implications in §4.

2. A Tidal Barrier for Accretion Flow

In this section we begin our analysis with an analytic approximation of the accretion flow pattern in two-dimensions. Since protoplanets accrete from a disk environment, it is informative to compare the disk structure with the gravitational potential of the protoplanet. The scale height in the direction normal to the plane of the disk is determined by the gas sound-speed c_s such that

$$H = c_s/\Omega, \quad (1)$$

where $\Omega = (GM_*/a^3)^{1/2}$ is the Keplerian frequency of the disk and M_* is the mass of the central star. The Bondi radius for a steady, spherically symmetric accretion flow onto a protoplanet with a mass M_p is

$$R_B = GM_p/2c_s^2. \quad (2)$$

The Roche radius for a planet with semi major axis a is

$$R_R = (M_p/3M_*)^{1/3}a. \quad (3)$$

The magnitude of

$$\frac{R_R}{R_B} = \frac{2H^2}{3R_R^2} = 2h_1^2 \left(\frac{1}{3q_3^2} \right)^{1/3} \quad (4)$$

and,

$$\frac{H}{R_B} = \frac{2h_1^3}{q_3} \quad (5)$$

where $q_3 \equiv M_p/10^{-3}M_*$ and $h_1 = c_s/(0.1\Omega a)$ are the normalized planetary mass and disk scale height. In a minimum mass nebula model (Hayashi et al. 1985) where the gas temperature is assumed to be identical to the equilibrium temperature of the large grains, $h_1 \simeq 0.7$ at the present radii of Jupiter ($a = 5.2$ AU).

Under these conditions, the critical stage where the accretion rate is being suppressed occurs when $R_R \sim R_B \sim H$ and $M_p = M_J$. In principle, this region should be analyzed with full 3D consideration (Bate et al. 2003; Klahr & Kley 2006, cf.). However, the inadequacy of present-day facilities and the enormous computational requirements of full 3D numerical simulations place severe limits on both numerical resolution and temporal range. It is therefore useful to first consider a more idealized 2D flow pattern. Through numerical simulations and analysis in this section, we emphasize that the protoplanets' accretion rate is sensitive to the global evolution of the disk structure. This approximation is also reasonable in the limit that $R_R > H$ or equivalently $R_B > R_R$.

There have been multiple simulations studying the process of accretion onto a protoplanet covering a wide range of approaches. In an attempt to achieve high resolution near the protoplanet, several investigators (Korycansky & Papaloizou 1996; Tanigawa & Watanabe 2002) considered flows in a local region surrounding the protoplanet. Both studies assume that gas entering the simulation from both the leading and trailing azimuthal boundaries has a constant surface density and Keplerian velocity. A second set of simulations have investigated the global disk response to protoplanet-disk interaction. Some of these simulations include the protoplanet’s orbital migration while others include the effect of intrinsic turbulence. In all cases, they show that Jupiter mass protoplanets induce gap formation in protostellar disks with properties similar to that of the minimum mass nebula. In order to achieve sufficient resolution for studying the flow around protoplanets, most of these simulations are carried out for M_J -mass protoplanets in disks with $H/a \sim 0.07$ such that their $R_R \sim R_B \sim H$. Here, we are primarily interested in the disk regions where $R_B \gg R_R$ which is possible only for $R_R > H$. We show that gas-giants with $M_p < M_J/3$ (such as Saturn) are probably formed in regions of the disk that satisfy $H/a < 0.04$ during the epoch of planet formation.

2.1. Streamlines Intersecting The Feeding Zone

The continuity and momentum equations for a two-dimensional inviscid disk can be obtained through vertical integration such that

$$\frac{\partial \Sigma}{\partial t} + \nabla \cdot (\Sigma \mathbf{v}) = 0 \quad (6)$$

and,

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} = -2\Omega \hat{\mathbf{z}} \times \mathbf{v} - \nabla \Phi_g - c_s^2 \nabla \ln \Sigma. \quad (7)$$

Similar to the 1D analysis presented in §2.2, we have adopted an isothermal equation of state. We determine the magnitude of $\mathbf{v}$ in a frame which rotates at the same angular frequency Ω as the planet orbits its host star. In the analytic derivations, we neglect the effect of viscous and shock dissipation, but some degree of dissipation is present in our numerical simulations. These approximations are adequate for linking streamlines between shock waves. We also adopt a Hill’s approximation for the potential in this frame such that

$$\Phi_g = -\frac{3}{2}\Omega^2 x^2 - \frac{GM_p}{(x^2 + y^2)^{1/2}} \quad (8)$$

where the shearing-sheet coordinates (Goldreich & Lynden-Bell 1965) are centered around the planet with x and y correspond to the radial and azimuthal directions. Although this

expression is a much more accurate description of Φ_g than the one-dimensional approximation given in Equation (19), approximations must be made to solve it analytically.

In the absence of any viscous or nonlinear dissipation, it is helpful to analyze the flow with a vortical stream function (ψ) and a scalar potential (ϕ) which varies along the lines of constant ψ such that the 2D mass flux can be decomposed into

$$\Sigma \mathbf{v} = \nabla \times \hat{\mathbf{z}} \psi + \nabla \phi. \quad (9)$$

Contours lines of constant ψ and ϕ are orthogonal to each other, and a steady-state ($\nabla \cdot (\Sigma \mathbf{v}) = 0$) corresponds to $\nabla^2 \phi = 0$. The velocity is parallel to streamlines of constant ψ , along which the mass flux (Σv) is conserved, and the momentum equation can be reduced to two integrals of motion

$$\frac{\omega + 2\Omega}{\Sigma} = \varpi(\psi) \quad (10)$$

$$v^2/2 + c_s^2 \ln \Sigma + \Phi_g = E_B(\psi) \quad (11)$$

which correspond to the vortensity (Papaloizou & Lin 1989) and Bernoulli constant (Korycansky & Papaloizou 1996; Balmforth & Korycansky 2001).

Differentiating the mass flux $\mathbf{F}_m \equiv \Sigma \mathbf{v}$ and E_B with respect to ϕ along these streamlines,

$$(d \ln \Sigma / d \phi)_\psi = -(d \ln v_\psi / d \phi)_\psi, \quad (12)$$

and,

$$\frac{(v_\psi^2 - c_s^2)}{\Sigma} \left(\frac{d \Sigma}{d \phi} \right)_\psi = - \left(\frac{\partial x}{\partial \phi} \right)_\psi \frac{\partial \Phi_g}{\partial x} - \left(\frac{\partial y}{\partial \phi} \right)_\psi \frac{\partial \Phi_g}{\partial y} \quad (13)$$

$$= \left(\frac{x}{\Sigma v_x} \right) \left(3\Omega^2 - \frac{GM_p}{(x^2 + y^2)^{3/2}} \right) - \left(\frac{y}{\Sigma v_y} \right) \frac{GM_p}{(x^2 + y^2)^{3/2}} \quad (14)$$

where we define the magnitude of $\mathbf{v}$ along the streamline to be v_ψ . Equation (14) clearly indicate a potential transonic point along each streamline.

2.2. Analogous Bondi Solution

The sonic transition in a 2D flow is similar to that in the 1D flow, and the quantities F_m and E_B are equivalent to the mass flux

$$\dot{M}_p = 4\pi \rho u r^2 \quad (15)$$

and the one-dimensional Bernoulli energy

$$E_{B1} = u^2/2 + c_s^2 \ln \rho + \Phi_g \quad (16)$$

where ρ and u are the spherically symmetric density and radial velocity. We find it is useful to identify the dominant physical effects which may affect the 2D flow pattern with the help of an approximate 1D solution.

In a steady state, the equations of 1D continuity and motion are combined into

$$\frac{(c_s^2 - u^2)}{u} \frac{du}{dr} = -\frac{2c_s^2}{r} - \nabla \Phi. \quad (17)$$

In the standard Bondi analysis for a point-mass potential in which $\Phi_g = \Phi_0 \equiv -GM_p/r$, the continuity of a steady flow across the sonic point r_s where $u = c_s$ requires $r_s = -2c_s^2/\nabla \Phi$ and $E_B = c_s^2 \ln \rho_\infty$. By scaling the background quantities with those at the sonic point r_s such that $\rho_s = \rho_\infty \exp 3/2 \sim 4.5\rho_\infty$, we find the accretion rate to be

$$\dot{M}_p = \dot{M}_{pm} \simeq 14\rho_\infty G^2 M_p^2 / c_s^3. \quad (18)$$

In this case, the potential at r_s is lower (more negative) than at $r = \infty$, and the gravitational focusing effect causes $\rho_s > \rho_\infty$.

However, around a protoplanet with a circular orbit and semi major axis a , the region outside its Roche radius ($R_R = (M_p/3M_*)^{1/3}a$) is dominated by the gravity of its host star. In the frame which co-rotates with its orbit, the full tidal potential due to both the planet and its host star is given by Equation (8). For the 1D analysis in this section, we introduce an approximation to the gravitational potential which includes the tidal effect induced by central star but neglects its gravitational torque on the otherwise assumed spherically symmetric (relative to the protoplanet) radial flow such that

$$\Phi_g = \Phi_0 \left(1 + \left(\frac{R_R^2}{r^2} + \frac{R_R^2}{rR_\infty} \right)^{-1} \right). \quad (19)$$

Here we did not include the Coriolis force associated with the rotation of the frame. In this approximation, gravity reduces to that of a point mass ($\nabla \Phi \simeq GM_p/r^2$) for $r < R_R$, it becomes repulsive, ($\nabla \Phi \simeq -GM_p/[R_R^2(1 + r/R_\infty)^2] + GM_p/r^2$) for $R_R < r < R_\infty$, and vanishes asymptotically ($\nabla \Phi \simeq -GM_p R_\infty^2 / R_R^2 r^2$) for $r \gg R_\infty$. In principle, this approximation cannot be used to replace the full multi-dimensional simulation on the detailed structure and dynamics of the outer most region of the envelope near the protoplanet's Roche radius. However, by setting $R_\infty = r_{\max} = \sqrt{12}R_R$ (Greenzweig & Lissauer 1990), the potential in the frame which co-rotates with the protoplanet's orbit can be reasonably approximated along the radial direction.

For the tidal potential, the right hand side of Equation (17) vanishes at $r_s \equiv R_R/\xi$ where the parameter ξ can be obtained from a quadratic equation

$$\xi^2 + \left(\frac{2R_R}{R_\infty} + \frac{R_R}{R_B} \right) \xi + \left(\frac{R_R^2}{R_\infty^2} + \frac{2R_R^2}{R_\infty R_B} - 1 \right) + \frac{R_R^3}{R_B R_\infty^2} \frac{1}{\xi} = 0 \quad (20)$$

the solution of which is plotted in Figure (1). In the limit that $R_R < R_B$, $\xi \simeq 1 - R_R/R_\infty + R_R/2R_B$, i.e. the sonic point is close to the Roche radius.




Fig. 1.— Sonic radius parameter ξ , defined by the sonic point $r_s = R_B/\xi$, as a function of the ratio of Roche to classical Bondi radii. The dependence of ξ on R_R/R_B can be found in Equation (20).

When the tidal potential is taken into account, $E_B = c_s^2 \ln \rho_\infty - GM_p R_\infty / R_R^2$ at large r , such that the density at the sonic point is given by

$$\rho_s = \rho_\infty \exp \left[\frac{\Phi(r_s) - \Phi(\infty)}{c_s^2} - \frac{1}{2} \right]. \quad (21)$$

Since the absolute value of the potential at r_s is near a maximum (where gravity $\nabla \Phi \sim 0$), $\rho_s < \rho_\infty$. This density decline is due to the need of a positive pressure gradient, required to overcome the tidal barrier associated with this potential maximum. In the limit of relatively small c_s (or equivalently large R_B) a large density reduction is needed. Based on the above extrapolation that $r_s \sim R_R$ for $R_R \ll R_B$,

$$\rho_s \simeq 0.6 \rho_\infty \exp(-\eta R_B / R_R) \quad (22)$$

where $\eta \equiv 2(R_\infty / R_R - 2) \sim 3$ is positive. In a tidal potential with $R_R \ll R_B$, $\xi \sim 1$, $r_s \sim R_R$, and

$$\dot{M}_p = \dot{M}_t \simeq \left(\frac{2\pi}{3^{1/3}} \right) c_s R_R^2 \rho_\infty \exp \left(-\eta \frac{R_B}{R_R} \right) = 0.135 \dot{M}_{pm} \left(\frac{R_R}{R_B} \right)^2 \exp \left(-\eta \frac{R_B}{R_R} \right). \quad (23)$$

The above equation indicates that, in the limit $R_R \ll R_B$, the tidal accretion rate ($\dot{M}_t$) can be much smaller than the standard Bondi accretion rate for a point mass potential $\dot{M}_{pm}$.

2.3. Boundary Conditions in 2D Flows: Loading the Streamlines

A comparison of the governing equations (17) and (14) indicates that the competing effects of pressure gradient and gravity in the 2D flows is determined by the boundary conditions of the streamlines. At large x distance from the planet, the unperturbed 2D flow is approximately Keplerian, with $v_x \sim 0$ and $v_y \sim -3\Omega x/2$, which can be substantially larger than the sound-speed. These streamlines are on hyperbolic paths with large impact parameters relative to the protoplanet and material on these streamlines will not accrete onto the planet.

The streamlines which are relevant to accretion flow are those with impact parameters smaller than R_R . In the limit $H \ll R_R$, pressure does not contribute significantly, and Equation (7) reduces to that for non-interacting particles. Particles that originate from unperturbed Keplerian orbits at $x_\infty = x_{\max} \equiv \sqrt{12} R_R$ and $y_\infty = \pm \pi a$, reach the vicinity of the unstable Lagrangian saddle points $(\pm R_R, 0)$ with negligible velocity (Greenzweig & Lissauer 1990). The region of the disk between $a - x_\infty$ and $a + x_\infty$ is commonly referred to as the feeding zone. Although particles with initial Keplerian orbits that pass through the location

$y_\infty = \pm\pi a$ with $x_\infty = 1.8x - 2.7R_R$ can directly strike a compact protoplanet core, particles' on neighboring orbits are strongly scattered by the planet (Bryden et al. 2000).

Although ϖ and E_B are useful quantities for analyzing the flow in the horseshoe region, they are not conserved for gas accreted from the disk onto the planet. More specifically, for fluid in the protostellar disk with Keplerian asymptotic velocities, $\varpi = \Omega/2\Sigma_\infty$. However, if gas enters into the planet's Roche radius with $x^2 + y^2 < 3R_R^2/4$ in a Keplerian circumplanetary disk, its streamlines have $\omega < -2\Omega$ and $\varpi < 0$. This mismatch in ϖ implies that gas can only be accreted from protostellar disks onto protoplanetary disks through shock dissipation (Tanigawa & Watanabe 2002, see 2D simulations by). Since any dissipation would invalidate the conservation of ϖ and E_B , the flow pattern derived from these conservative quantities cannot adequately describe accretion flow onto the protoplanet (Korycansky & Papaloizou 1996, cf.).

However, these quantities are useful for determining the conditions under which gas on some streamlines in the horseshoe regions may avoid shock dissipation and accrete onto the protoplanet. The flow in this region depends sensitively on the boundary condition at (x_∞, y_∞) . If the flow at (x_∞, y_∞) is strictly Keplerian, as assumed in some previous local simulations, $E_B = c_s^2 \ln \Sigma_\infty - 3\Omega^2 x_\infty^2/8$. In this case,

$$\Sigma_s \equiv \Sigma(x_s, y_s) = \Sigma_\infty \exp \left[-\beta \frac{R_R^2}{H^2} - \frac{1}{2} \right] \quad (24)$$

at the sonic point (x_s, y_s) where $u = c_s$. The parameter β is independent of H or c_s . For Keplerian boundary conditions, the value of

$$\beta = \frac{3}{2} \left[\left(\frac{x_\infty^2}{4R_R^2} \right) - \frac{x_s^2}{R_R^2} - \frac{2R_R}{(x_s^2 + y_s)^{1/2}} \right]. \quad (25)$$

In this expression for β , the actual values of (x_s, y_s) can only be determined by the detailed solutions of Equations (6) and (7).

In analogy with the 1D Bondi solutions, we suggest that there may exist a sonic transition point in Equation (14) as is found in Equation (17). In this case, if $|u| < c_s$ at some point along the streamline, the right hand side of Equation (14) would vanish during the sonic transition at (x_s, y_s) with

$$\frac{v_x(x_s, y_s)}{v_y(x_s, y_s)} = \frac{x_s}{y_s} \left(\frac{(x_s^2 + y_s^2)^{3/2}}{R_R^3} - 1 \right). \quad (26)$$

The point (x_s, y_s) is close to the turn-around point (where u_y vanishes) of the horseshoe

orbit for non-interacting particles. Since $v_x^2 + v_y^2 = c_s^2$ at (x_s, y_s) , we can express,

$$\frac{v_y^2}{c_s^2} = \left[1 + \left(\frac{(x_s^2 + y_s^2)^{3/2}}{R_R^3} - 1 \right)^2 \left(\frac{x_s}{y_s} \right)^2 \right]^{-1}. \quad (27)$$

In the 1D approximations (see the previous subsection), the physical parameters at the sonic points can be analytically determined from those at infinity. But in 2D flows, Equations (14) and (27) generally do not lead to an analytic relation between r_s and c_s . However, their approximate solutions suggest that the condition for $v_x^2 < c_s^2$ and $v_y^2 < c_s^2$ is $r_s^2 \equiv x_s^2 + y_s^2 \sim R_R^2$, *i.e.* in the 2D limit, the transonic radius is also comparable the planet's Roche radius ($r_s \sim R_R$) and

$$\dot{M}_p \simeq 2\pi R_R c_s \Sigma_s = 2\pi R_R H \Omega \Sigma_\infty \exp \left[-\beta \left(\frac{R_R^2}{H^2} \right) - \frac{1}{2} \right]. \quad (28)$$

The above expression clearly indicates the importance of β . If $\beta < 0$, $\dot{M}$ will decrease with H as shown in the local simulations of Tanigawa & Watanabe (2002). However, if $\beta > 0$, $\dot{M}$ will increase with H as shown in the global simulation by Bryden et al. (1999).

We suggest this dichotomy can be attributed to the boundary conditions at (x_∞, y_∞) . In order to illustrate this conjecture, we first consider a special streamline which passes through a sonic transition at the same azimuthal phase as the planet (*i.e.* $y = 0$), the regularity condition for Equation (14) requires $\partial\Phi_g/\partial x = \partial\Phi_g/\partial y = 0$ or equivalently, $x_s = R_R$. In the limit that $H \ll R_R$, $\Phi_g(x_s, y_s) = -9\Omega^2 R_R^2/2$, $x_\infty = x_{\max} = \sqrt{12}R_R$, $y_\infty = \pi a$, and $\beta \sim 0$. These streamlines are analogous to the parabolic trajectories of those particles which can marginally reach the separatrix at the Lagrangian points from an initial circular Keplerian orbit. Starting with Keplerian motion, streamlines with $x_\infty > x_{\max}$ do not enter into the planet's Roche lobe. However, Keplerian streamlines with $R_R < x_\infty < x_{\max}$ can enter into the planet's R_R and cross the sonic radius with very small negative values of β . In local simulations done previously, the disk flow at some specified boundaries is explicitly set to be Keplerian at an azimuthal phase $y_\infty < \pi a$. This artificial choice of boundary condition leads a small negative value of β for disks with both negligible and finite H 's. This result implies that tidal barrier is overcome purely by the initial kinetic energy rather than the positive pressure gradient and Σ_s is slightly larger than Σ_∞ as a consequence of convergent flow. If β is negative, Equation (28) indicates that $\dot{M}_p$ is a decreasing function of H . This is in agreement with the simulations of Tanigawa & Watanabe (2002). With their value of Σ_∞ ($\sim \Sigma_{mn}$), the growth timescale would be $\tau_p = M_p/\dot{M}_p \sim 10^3 \text{yr} (M_p/10M_\oplus)^{-0.3}$, which is much shorter than the global depletion timescale (τ_{dep}) for the disk. The discrepancy between these timescales illustrates the need for a mechanism to quench the accretion flow.

In contrast to local simulations, full-scale 2D simulations the protoplanet induces the formation of a gap in the disk through both accretion onto itself and tidal dissipation within the disk (Bryden et al. 1999). In such simulations the boundary conditions at (x_∞, y_∞) are significantly different from those assumed in local simulations. Near the edge of the gap, the perturbed pressure gradients cause the velocity of the gas to become super Keplerian at positive x_∞ and sub Keplerian at negative x_∞ . The magnitude of the flow velocity,

$$v_y(x_\infty, y_\infty) = \Omega a \left(\frac{H^2}{2a^2} \frac{\partial \ln \Sigma}{\partial \ln x} - \frac{3x}{2a} \right)_{x=x_\infty}, \quad (29)$$

is reduced for both positive and negative value of x_∞ . The typical gap has a width $R_R < \Delta < x_{\max}$. When the thermal and viscous conditions for gap formation ($R_R > H$ and $M_p/M_* > 40\nu/\Omega a^2$ where ν is the effective viscosity) are satisfied, the surface density inside the gap $\Sigma_\infty(x_\infty, y_\infty) \sim \epsilon \Sigma_o$ with $\epsilon < 10^{-2} - 10^{-4}$ (Bryden et al. 1999). Although this reduction in Σ_∞ decreases the value of $\dot{M}_p$ (by a factor similar to ϵ), the growth timescale τ_p would still be more than an order of magnitude shorter than τ_{dep} if the protoplanet is able to accrete all the gas which diffuses into the gap region either through turbulence (Nelson & Papaloizou 2003) or due to intrinsic or artificial viscosity (Artymowicz & Lubow 1994; Kley et al. 2001).

In the limit of severe surface density depletion near the protoplanet, the magnitude of v_y within the gap may be substantially reduced from the Keplerian approximation. In the limit $|x_\infty| < H$, we approximate Equation (29) as $v_y^2(x_\infty, y_\infty) = (9/4)\Omega^2 x_\infty^2 (1 - f_{\text{sub}})$ where

$$f_{\text{sub}} = \frac{2}{3} \frac{H}{x} \frac{H}{a} \frac{\partial \ln \Sigma}{\partial \ln x}. \quad (30)$$

For values of $0 < f < 1$, the Bernoulli constant can be written as $E_B \simeq c_s^2 \ln \Sigma_\infty - 3(1 + 3f_{\text{sub}})\Omega^2 x_\infty^2/8$. At large azimuthal phases, the streamlines (constant ψ) are nearly parallel to the y -axis. Along these streamlines, the potential has a maximum at $\partial \Phi_g / \partial x = 0$, which is located on a circle $x^2 + y^2 = R_R^2$ centered on the protoplanet. In order for the background gas to reach the protoplanet within the Roche radius, it must pass through this local potential maximum with a positive pressure gradient. In the limit of low sound-speed, a sufficiently strong pressure gradient is only attainable with a large density gradient, *i.e.* the severe gas depletion interior to R_R . For these conditions, the value of

$$\beta = 3 \left[\frac{1}{8} \left(\frac{(1 + 3f_{\text{sub}})x_\infty^2 - 4x_s^2}{R_R^2} \right) - 1 \right]. \quad (31)$$

In the limit $x_\infty \simeq x_{\max}$ and $x_s \sim R_R$, $\beta \simeq 27f_{\text{sub}}/2 > 0$. For any value of $0 < f_{\text{sub}} < 1$, $\beta > 0$ for all streamlines with $x_\infty > (\sqrt{12}/(1 + 3f_{\text{sub}})) R_R$. Streamlines with smaller values of x_∞ attain horseshoe orbits that do not enter into the Roche radius. In cold disks ($H < R_R$) this

tidal barrier suppresses the flow in the azimuthal direction such that only a small fraction of the gas which is diffused into the gap can actually reach its Roche lobe. Consequently, the accretion rate derived from Equation (28) become substantially smaller than that obtained when neglecting the effect of the azimuthal tidal barrier.

2.4. Protoplanet’s Asymptotic Mass

We now apply the appropriate boundary condition in protostellar disks to determine the asymptotic mass of protoplanets. For illustration simplicity, we first adopt the 1D approximation and then consider the full 2D flow.

The density of the ambient gas can be derived from $\rho_\infty = \Sigma/2H$ where the surface density is $\Sigma = f_g \Sigma_{mn}$ and f_g is a scaling factor. We use the minimum mass nebula model $\Sigma_{mn} \sim \Sigma_0(a/1\text{AU})^{-3/2}$, where $\Sigma_0(t_0) = 2 \times 10^3 g/cm^3$ as a fiducial prescription. We also approximate the evolution of the disk by $\Sigma = \Sigma_0 \exp(-t/\tau_{\text{dep}})$ (Ida & Lin 2004) where the characteristic gas depletion timescale τ_{dep} inferred from the observed SED is $\tau_{\text{dep}} \sim 3 - 10 Myr$.

From the 1D mass accretion rate given in Equation (23), we can determine the asymptotic mass of the planets M_f by first solving for $x_f = 3^{1/6}(\eta/2)^{1/2}h_1^{-1}q_{3f}^{1/3}$ with

$$\int_0^{x_f} \exp x^2 dx = A_0 \frac{f_g \Sigma_{mn} a^2}{h_1 M_*} \frac{\tau_{\text{dep}}}{P} \quad (32)$$

where $A_0 = (10^8 2^3/3)^{1/2} \pi^2 \eta^{1/2}$ is a constant, and the planet’s asymptotic mass is contained in $q_{3f} \equiv M_f/10^{-3} M_*$. The left hand side of Equation (32) is a rapid increasing function of x_f . For a minimum mass nebula model (*i.e.* $f_g = 1$ at $a=5$ AU) the magnitude of its right-hand side is $\sim 10^6$ and the solution of Equation (32) is $x_f \sim 4$. However, the protoplanet may be embedded inside a gap where the gas density is reduced by at least 2 orders of magnitude from that in the minimum mass nebula. Long period protoplanetary embryos may also have formed during the advanced stages of disk evolution when the magnitude of f_g may also be relatively small. However, the value for x_f is relatively insensitive to the magnitude of f_g and it reduces to ~ 3 if $f_g \sim 10^{-2}$. The corresponding value for the asymptotic planet mass is

$$M_f \simeq \frac{x_f^3}{3^{1/2}} \left(\frac{2}{\eta}\right)^{3/2} \left(\frac{H}{a}\right)^3 M_*. \quad (33)$$

The structural parameters of the disk are functions of M_* and a . Near the snow line of a minimum mass nebula, the magnitude of M_f is comparable to that of Jupiter. Equation (33)

is also equivalent to $R_R(M_f) \sim H$. This relation is in agreement with the thermal condition for the tidal truncation and gap formation of protostellar disks by embedded protoplanets (Lin & Papaloizou 1986).

The dependence of Σ_s/Σ_∞ on the sign of β in 2D is similar to that of ρ_s/ρ_∞ on the presence of the tidal barrier in the 1D approximation. Using the 2D mass accretion rate given by Equation (28), together with the approximate global Σ depletion formula, we deduce an asymptotic mass for a protoplanet of

$$\frac{M_p}{M_*} = \frac{3}{|\beta|^{3/2}} \left(\frac{H}{a} \right)^3 \left[\ln \frac{8\pi|\beta|}{9} \left(\frac{aM_{\text{gap}}\tau_{\text{dep}}}{HM_*P} \right) \right]^{3/2} \quad (34)$$

where $M_{\text{gap}} = \pi\epsilon\Sigma_o a^2$ is the characteristic mass associate with the gap region. For $\beta \sim 1$ and $\epsilon \sim 10^{-3}$, the above asymptotic mass would yield $R_R \sim H$. However, if the magnitude of β is very small or negative, the above equation reduces to $M_p/M_* \sim 3(8\pi HM_{\text{gap}}\tau_{\text{dep}}/9\pi aM_*P)^{3/2}$, which is substantially larger. To calculate the actual values of β and f_{sub} , we must utilize numerical simulations

3. Two Dimensional Numerical Simulations

3.1. Numerical Method

In order to verify the 2D analytic approximation in the previous section and deduce a value for β , we now present a series of 2D numerical simulations. The 2D numerical scheme we use is a fully parallel hydrodynamical code with which the continuity and momentum equations are solved on a fixed Eulerian grid in 2D cylindrical coordinates (r, ϕ) . A staggered grid is introduced where scalars are defined at grid centers and vectors are defined on grid edges, allowing us to maintain second-order spatial accuracy. Earlier versions of the code were used, among other applications, to study the formation of gaps in protostellar disks (Bryden et al. 2000) and also boundary layer physics in the accretion disks around cataclysmic binaries (Kley 1989). The equations of continuity and motion for the fluid and the expression for the gravitational potential are similar to those given in Equations (6), (7), and (8) respectively. In the equation of motion, we adopt an isothermal equation of state $P = c_s^2 \rho$, with a radially varying sound-speed, $c_s = (H/a)V_{\text{kep}}$ where the aspect ratio of the disk H/a is assumed to remain constant. In order to make a direct comparison with the results of the analytic approximation, we neglect the effect of the viscous stress associated with the turbulent motion. A softening parameter $r_{\text{soft}} = 0.5R_R$ is adopted for the gravitational potential of the planet.

The planet is initialized with a $1M_{jup}$ planet at a location $(r, \phi) = (1AU, 0)$. Although the planet is held fixed, any mass accreted during the simulation adds to the overall gravitational potential of the planet. The simulations cover the entire azimuthal extent of the disk and a radial range from $0.5AU$ to $1.5AU$. Simulations were initialized with a surface density profile of $\Sigma = 950 \frac{g}{cm^3} \left(\frac{r}{1AU}\right)^{-3/2}$, and have a numerical resolution of $(r, \phi) = (200, 1600)$.

Each of the terms in the fluid equations are dealt with consecutively through both operator and directional splitting. Our advection scheme is an extension of the simple first-order upwind scheme based on the prescription of van Leer (1977), and allows for the detailed treatment of strong shocks. Details of the advection scheme can be found in Kley & Hensler (1987). We solve the entire problem in a frame rotating with the planet.

3.2. Two Dimensional Numerical Models

Before the fluid is incorporated onto the planet it must navigate through both the shocks and sub-disk surrounding the protoplanet. In the 2D study we neglect any radiative feedback onto the gaseous disk from the energy released during the cooling and contraction of the planetary atmosphere. Following Tanigawa & Watanabe (2002), we decrease the surface density of the disk in the region $r_{sink} \leq 0.5R_R$ at a rate given by

$$\Sigma(t + \Delta t) = \Sigma(t) \left[1 - \frac{\Delta t}{\tau_{sink}} \right]. \quad (35)$$

This mass is added to the mass of the protoplanet, and gradually increases its gravitational contribution. Tanigawa & Watanabe (2002) demonstrated that the overall accretion rate ($\dot{M}$) was relatively unaffected by the choice of τ_{sink} provided that $r_{sink} \approx 0.1R_R$. Because of our lower numerical resolution near the protoplanet we set $r_{sink} = 0.5R_R$ and $\tau_{sink} = 1/\Omega$. The dominate effect which regulates the accretion rate is the supply of material to the sub-disk region. This quantity is determined by the effectiveness of loading the streamlines far from the planet, so the values of r_{sink} and τ_{sink} should not be crucial in determining the asymptotic mass of the planets. In order to test the validity of this assumption and the convergence of our results, we carried out several simulations with varying values of τ_{sink} , but these showed little variation in the overall accretion rate.

We present the results of two models in which $H/a = 0.04$ and 0.07 respectively. Figure (2) shows both the total mass and accretion timescale as a function of orbital period for these models. The accretion rates approach a steady state on the time scale of 100 orbital periods. This time scale is comparable to the liberation period of flows in the horseshoe region and the synodic period of the flow near the separatrix at the Lagrangian points. It reflects the

protracted adjustment of the disk structure to the tidal perturbation of the protoplanet. The accretion timescale $(\tau_{1\text{AU}} = M_p/\dot{M}_p)_{1\text{AU}}$ for a Jupiter-mass protoplanet is proportional to both the local surface density ($\propto a^{3/2}$) and to its orbital period ($\propto a^{3/2}$). It can therefore be scaled to other radii such that

$$\tau_{\text{growth}} = \tau_{1\text{AU}}(a/1\text{AU})^3. \quad (36)$$

Figure (2) shows timescales for planet located at both 1 and 5AU. For a planet at 5AU, we see that the growth timescale becomes comparable to the gas depletion time scale of the disk. Within the computational duration, the total fractional gain in the planetary mass is 0.05 and 0.2 for $H/a = 0.04$ and 0.07 at 1 AU and one order of magnitude smaller at 5 AU. Protracted 2D simulations for protoplanets' nonlinear growth (over 10^4 orbits) require computational resources beyond that currently available.

The results in Figure (2) also clearly indicate that the mass of a planet embedded in a disk with a lower aspect ratio levels off at significantly lower mass than one embedded in a disk with larger (H/a) . Based on these results, we compute, from Equation (28), the value of

$$\beta \simeq \left(\frac{R_R^2}{H_2^2} - \frac{R_R^2}{H_1^2} \right)^{-1} \ln \left(\frac{\dot{M}_1 H_2}{\dot{M}_2 H_1} \right) \quad (37)$$

where the subscripts 1, 2 refer to the values for the two models. The value of β increases from a very small value to the order of unity after ~ 50 orbits (see Figure (3)). The disk is initialized as a Keplerian disk, and as a result there is some amount of mass on Keplerian orbits within the separatrix region ($R_R < x < \sqrt{12}R_R$) at azimuthal phase $y_\infty < \pi a$. This gas passes over the tidal barrier near the planet solely based on this artificial kinetic energy. For this reason the first ~ 30 orbits in Figure (2) show similar accretion rates onto the planets in both simulations and the initial value of β is more closely approximated by Equation (25). In fact, during this relaxation phase, the lower sound-speed simulation is able to accrete mass onto the planet faster because the outward pressure gradient that develops around the planet is smaller.

After approximately one libration period around the L_4 and L_5 points, mass elements that start their trajectory at the superior conjunction (*i.e.* $y = a\pi$) can reach the vicinity of the protoplanet's Roche radius. Thereafter, a large pressure gradient along the stream line is needed to overcome the tidal potential barrier. During this second phase, the accretion timescale can be up to an order of magnitude larger in the high sound-speed case. The value of $\beta \sim 1$ is more closely approximated by Equation (31). The discussion near Equation (29) indicates that this approximation is based on the assumption that $v_y(x_\infty, y_\infty) \simeq 0$. In Figure (4), we plot the value of v_y at the planet's superior conjunction as a function of the disk

radius. We note that within the feeding zone, the magnitude of v_y is significantly reduced from its Keplerian values which is in agreement with Equation (29).




Fig. 2.— The total mass of the planet as a function of time (top). The solid line shows results for $(\frac{H}{a}) = 0.04$, while the dotted line shows $(\frac{H}{a}) = 0.07$. The lower panel shows the accretion timescale for both simulations. The left-hand ordinate shows the timescale for a planet at 1 AU, while the right-hand ordinate shows the timescale for a planet located at 5 AU.




Fig. 3.— The evolution of β , deduced from the comparison of two models with $(\frac{H}{a}) = 0.04$ and 0.07 . From Equation (28), we see that a positive β -value indicates that $\dot{M}$ is an increasing function of the disk scale height (H).




Fig. 4.— The radial distribution of v_y at the protoplanet’s superior conjunction for models with $(\frac{H}{a}) = 0.04$ (solid line) and 0.07 (dashed line). A decline in the magnitude of v_y within the feeding zone is due to a pressure gradient associated with the surface density distribution across the gap.

Our simulations do not include explicit viscosity, and as a result mass diffusion into the gap is due to numerical viscosity or shocks. Figure (5) shows the total mass within the separatrix region ($|r - a| \leq \sqrt{12}R_R$) for the two simulations. Again, during the initial stages of the simulation the two cases track each other. The drop in gap mass is quite precipitous ($\sim 0.3M_{Jup}$) during the first 100 orbits. Comparison with Figure (2) shows that a majority of this gas is not accreted by the planet, but rather shocked by the planet and pushed out of the region. However, the lack of material is not the cause for the mass of the protoplanet to level off in the low sound-speed model; by the end of the simulation, there is more mass within the gap region in this simulation than in the high sound-speed simulation.




Fig. 5.— The total mass of the planet in the gap region ($|r - a| \leq \sqrt{12}R_R$) as a function of orbits for $\frac{H}{r} = 0.04$ (solid line) and $\frac{H}{r} = 0.07$ (dashed line). By the end of the simulation there is more mass within the gap for the low sound-speed case, indicating that the tidal barrier, not the lack of material within the feeding zone, is the cause of the planets increasing growth timescale.

In order to further illustrate that the pressure gradient is insufficient to overcome the tidal potential barrier along the stream line in the low sound-speed simulation, we identify streamlines with contours of constant vortensity ϖ . Since shocks occur only relatively close to the protoplanet, vortensity is approximately conserved along the stream lines in regions outside the Roche lobe. In Figures (6) and (7) we overlay the contours of ϖ on the velocity field. Outside R_R , the streamlines are nearly parallel, and those that eventually enter the protoplanet’s Roche lobe are initially located at $\sim 2 - 3R_R \sim 0.15 - 0.2a$ from its orbit.

Having determined that the flow is primarily along contours of constant ϖ , we can utilize the discussion at the end of §2.3 to link the flow velocity at the superior conjunction to the resulting accretion rate. In Figures (8) and (9), we plot $\Delta = (v_y + \Omega a)/v_{\text{kep}} - 1$, where $v_{\text{kep}} = (GM_*/a)^{1/2}$ and v_y is given in the rotating frame. Comparing to Equation (30), we see that $f_{\text{sub}} \simeq 2|\Delta|$. At $a = 0.8\text{AU}$ and 1.25AU (which correspond to $\sim \pm x_{\text{max}}$ away from the protoplanet), $|\Delta| \sim 0.03$ and thus $f_{\text{sub}} \sim 0.06$. For a streamline that connects $x_\infty = x_{\text{max}}$ and $x_s = R_R$, Equation (31) gives $\beta \sim 0.8 - 1.2$, which is in agreement with the results shown in Figure (3).




Fig. 6.— Potential vorticity (ϖ) contours overlying the velocity field of the disk for the low sound-speed simulation with $\left(\frac{H}{a}\right) = 0.04$. Potential vorticity, conserved in regions outside R_R , provides a good tracer of the streamlines within the flow.




Fig. 7.— The same as Figure (6), but for $\left(\frac{H}{a}\right) = 0.07$.




Fig. 8.— $\Delta = (v_y + \Omega a)/v_{\text{kep}} - 1$ at the protoplanet's superior conjunction for $(\frac{H}{a}) = 0.04$. This quantity parameterizes the deviation of the flow from Keplerian ($\Delta = 0$). Δ is related to $f_{\text{sub}} \simeq 2|\Delta|$, and can be used to determine the magnitude and sign of β , the accretion rate parameter.




Fig. 9.— Same as Figure (8) but for $(\frac{H}{a}) = 0.07$.

In Figures (10) and (11), we plot the variations of Σ along the azimuth at constant semimajor axis. Although the approximation of Equation (14) breaks down near the protoplanet, it provides a reasonable indication of the variation of Σ along the streamlines. The results in Figures (10) and (11) show much larger azimuthal amplitude variations in Σ for the low sound-speed model than for the high sound-speed model. The magnitude of Σ decreases by two orders of magnitude in $H/a = 0.04$ model. This result is consistent with our analytic discussion that a large density gradient is needed in the low sound-speed limit in order to sustain sufficient pressure gradient to overcome the tidal potential barrier, and can be compared to the 1D results of Figure (15).




Fig. 10.— Surface density (Σ) as a function of azimuthal angle (ϕ) at several values of constant r around the planet for $(\frac{H}{a}) = 0.04$ at $t = 500$ orbits (left panel). Contours are at $r = 0.95, 1.0$, and 1.05 AU, while the planet is located at 1 AU. The precipitous drop in density along azimuth provides the necessary pressure gradient force needed to overcome the tidal barrier. The right-hand panel shows the overall surface density in the region.




Fig. 11.— Same as Figure (10) but for $(\frac{H}{a}) = 0.07$. Given the higher sound speed, the density drop needed to overcome the tidal barrier is not as large as in the low sound-speed simulation.

Figures (12) and (13) show the Mach number $((v_x^2 + v_y^2)^{1/2}/c_s)$ distribution in the rotating frame near the planet for $(H/a) = 0.04$ and 0.07 respectively. Both the approximately Keplerian flow far from the planet, and the horseshoe orbits within the feeding zone are clearly visible in both plots. As observed in Tanigawa & Watanabe (2002), the radial extent of the shocks from the planet is proportional to the soundspeed. For the most part, the subsonic region is bounded by a strong shock. However, for the high sound speed case, the subsonic region also extends from inside the Roche lobe to the feeding zone beyond r_R , allowing gas to efficiently flow into the vicinity of the protoplanet. For the low sound-speed model, The subsonic region near the protoplanet is well separated from the low-velocity horseshoe region. These figures once again support the conjecture that large density gradient is needed for the pressure to overcome the tidal potential barrier.




Fig. 12.— The Mach number for $\left(\frac{H}{a}\right) = 0.04$ in a frame rotating with the planet. The dashed circle over-plotted on the flow denotes the approximate Hill sphere. The subsonic region surrounding the planet is well separated from the rest of the flow, and a large density gradient is needed to overcome the tidal barrier.




Fig. 13.— The same as Figure (12) but for $\left(\frac{H}{a}\right) = 0.07$. The subsonic region surrounding the planet is connected, via subsonic streamlines, to the material in the feeding zone. Thus, the density gradient needed to overcome the tidal barrier is not as large as in the $\left(\frac{H}{a}\right) = 0.04$ simulation.

3.3. Simulations Based on the 1D Tidal Barrier Approximation

The main objective in our simulation is to determine the asymptotic mass of protoplanets in depleting protostellar disks. The results of our analysis and numerical simulations (Figure (2)) indicates that protoplanets reach their asymptotic mass when their growth timescale becomes comparable to that of the disk depletion timescale, which is 10^5 time longer than their orbital period. The accreted gas collapses to form the envelope of gas-giant planets with size scales several times the present-day radius of Jupiter. The dynamical timescale near the core is $\sim 10^{-5}$ that of the orbital period. While we would like to simultaneously study the evolution of the gaseous envelope and the decline of the mass influx through the protoplanet’s Roche lobe, the limitation of currently available computational facilities prohibits the high-resolution, multi-dimensional simulations of the planetary accretion process over such a large range in the relevant timescales.

Here we carry out a series of 1D simulations based on the spherically symmetric approximation of the tidal potential (§2.2). The basic equations for the 1D code include three conservation laws of momentum, mass and energy (including radiative diffusion and convection), and the equation of state. During the phase of dynamical accretion, conventional numerical schemes based on a hydrostatic approximation (Pollack et al. 1996) are no longer adequate for the determination of the planetary structure and evolution. We developed a numerical scheme which is based on a Lagrangian approach where the radial coordinate evolves in time. It was developed to study the envelope structure as it undergoes a transition from quasi hydrostatic equilibrium to dynamical collapse (see a detailed description in (Li & Lin 2006)). The advantage of this method is that we can use it to resolve the large spatial range of the protoplanet’s envelope and to follow its evolution over an extended duration of time. The main limitations of this method are the assumption of spherical symmetry and our neglect of the protoplanet’s tidal torque on the global structure of the disk gas. We utilize the versatility of this scheme to highlight the dominant physical effects and compare our results to the 2D scheme to assess the validity of our 1D calculation at various stages of protoplanetary growth.

For the present application, gas is accreted onto a Saturn-mass object ($M_p = 100M_\oplus$). During the dynamic gas accretion stage of planetary formation, the envelope of the protoplanet will contract much more rapidly than before. While the gas flow close to the core remains in quasi-equilibrium, the outer region of the envelope, especially the region near the surface of the proto-planet, should experience dynamic gas accretion. We focused our calculations on the dynamic region of the gas flow, which covers the radial region from 10 core radii to 2 Bondi radii. Across the inner boundary, we calculated the gas accretion rate and infer a luminosity based on the assumption that the inflow is halted at the core radius. As

the gas flow in this region tends to a steady-state (to be shown below), the radiative transfer is very efficient and there is insufficient thermal pressure to counter balance the core’s gravity throughout this region. It is therefore a reasonable assumption for the velocity across the inner boundary to be close to the free-fall velocity. Constant temperature and density are set at the outer boundary to denote the nebula conditions. In order to approximate the effects of the tidal potential, we added a simple prescribed outward gravitational force in the equation of motion (Equation 19). We can then adjust the location of the Roche radius where gravity vanishes in order to assess the effects of this tidal barrier.

For comparison, we first constructed a test model for standard Bondi accretion with the nebula conditions of $T_{neb} \sim 50K$ and $\rho_{neb} \sim 1 \times 10^{-14}g/cm^3$ (Model 0 in Table 1). These parameters are relevant in the outer ($> 5 - 10$ AU) regions after a substantial fraction ($\sim 10^{-3} - 10^{-2}$) of the minimum mass nebula has already been depleted through global depletion or gap formation. In this, we neglect the radiative energy transfer and tidal barrier effects so that we can compare our numerical results with the analytic Bondi solution. We find a gas accretion rate of around $3 \times 10^{17}g/s$ from our numerical calculations, which is in agreement with the analytic Bondi solution $\dot{M} \simeq 1.4 \times 10^{11}(\frac{M}{M_{\odot}})^2(\frac{\rho(\infty)}{10^{-24}})(\frac{c_s(\infty)}{10km/s})^{-3}g/s$ (Frank et al. 1992).

We constructed another three models to investigate the tidal barrier effects, with different parameters listed in Table 1 (Model 1, 2, 3). In the table, T_{neb} denotes the temperature of the nebula gas, which corresponds to the sound-speed at the outer boundary of the envelope. The third column shows the ratio of Roche radius to the Bondi radius. The last column lists the average gas accretion rate for each model. The outer boundary condition for density is $1 \times 10^{-14}g/cm^3$ for all the models.

Model No.	T_{neb} (K)	$R_{\text{R}}/R_{\text{B}}$	$\dot{M}$ (g/s)
0	50	-	3.0×10^{17}
1	50	0.5	1.6×10^{14}
2	50	1.0	1.7×10^{17}
3	100	1.0	6.2×10^{16}

Table 1: Parameters for the single, one-dimensional models. Model 0 shows Bondi accretion, which does not include radiative energy transfer or the tidal barrier. T_{neb} denotes the temperature of the nebula gas, while $\dot{M}$ is the average gas accretion rate calculated from the model.

We evolved all the models to steady-state, where the accretion rate became constant with radius. Figure (14) shows the temperature profiles of the three models after they have reached steady-state. As we can see from the figure, the outer regions of the envelopes are optically thin, cool, and isothermal for all the models. Figure (15) plots the density profiles of the three models. The model with $R_R/R_B \sim 0.5$ shows a much larger density gradient and thus much smaller density at the location of Roche radius than the other two models. At small radii, the pressure gradient does not significantly perturb the nearly free-falling motion of the gas. However, as the stellar gravity acts to repel the gas outside the Roche radius, sufficiently large positive pressure and density gradients are needed to overcome the tidal barrier. Therefore, in the case of $R_B > R_R$, this gradient leads to a density at R_R that is much smaller than that in the disk. Since free fall onto the core can only proceed inside R_R and the flow across R_R is limited by the speed of sound, the mass accretion rate is much reduced from the conventional Bondi formula. Figure (16) shows the evolution of gas accretion rate with time at the locations of Roche radius for three models. Nearly constant accretion rate indicates that all the models have reached steady state after approximately $6 \times 10^9 s$. Since the flow between R_B and R_R is significantly perturbed by the gas pressure gradient and softened gravity, this time scale must be at least several times the local dynamical free-fall time scale ($\sim 5 - 8 \times 10^8 s$). However, in contrast to the 2D simulations, the flow does not need to adjust over the liberation period of flows in the horseshoe region and the synodic period of the flow near the separatrix at the Lagrangian points.

From the two models with the same R_R/R_B but different nebula temperature (Model 2 and Model 3), it's found that higher nebula temperature leads to slightly lower gas accretion rate. This behavior is similar to what we find in the Bondi solution, in which $\dot{M} \propto c_s^{-3}(\infty)$. When comparing the models with the same nebula temperature but different R_R/R_B (Model 1 and Model 2), we find that the accretion rate is reduced by three orders of magnitudes for the model with $R_B > R_R$. Thus, the tidal barrier significantly suppresses in the gas accretion across the Roche radius. Even if the models are surrounded by nebula gas with the same density, only a small fraction of that gas can be accreted by the model with $R_B > R_R$.




Fig. 14.— Temperature profiles as a function of radius for 1D models 1, 2, and 3 after they have reached steady-state. The parameters for each model are listed in Table (1).




Fig. 15.— Density profiles as a function of radius for the three 1D models after they have reached steady-state. The denotations are the same as in Figure (14).




Fig. 16.— Gas accretion rate as function of time for the three models. Different line types correspond to the different models listed in Table (1).

We further compute two series of disk models (4 and 5) for planetary masses ranging from $30M_{\oplus}$ to 100 and $300M_{\oplus}$, respectively. Models in series 4 have (T, ρ_{neb}) of $(100\text{K}, 10^{-12}\text{g cm}^{-3})$, while models in series 5 have (T, ρ_{neb}) of $(50\text{K}, 10^{-12}\text{g cm}^{-3})$. We determine the magnitude of $\dot{M}$, as a function of M_p after it has attained a steady state value at 10 AU (see Figure 17). These results are in good agreement with the relationship in Equation (23). Based on these results, we determine the total mass of the protoplanet after 20 Myr of accretion onto a core with an initial mass of $30M_{\oplus}$ (see Figure 18). In both model series we neglect the decline of gas density associated with formation of the gap in the disk, but include that due to the global depletion over a time scale $\tau_{\text{dep}} = 3\text{Myr}$. Despite these simplifications, the asymptotic values of M_p (100 and $300 M_{\oplus}$ for $T_{neb} = 50\text{K}$ and 100K respectively) are comparable to that found in Equation (33) with the appropriate boundary conditions.

4. Summary and Discussion

In this paper, we present evidence to suggest that the asymptotic mass of protoplanets is determined by the structure of their nascent disk. The main physical process involved here is that the tidal potential of the central star provides a barrier to the gas flow in the vicinity of the protoplanet. Through its tidal interaction with the disk, a growing protoplanet exerts an increasingly strong torque which first leads to the formation of a gap. Although the disk gas may continue to diffuse into the gap, it is forced to flow along the horseshoe stream lines. In the absence of any dissipation, the vortensity of the flow along the stream lines is conserved. But near the protoplanets, shock dissipation can lead to vortensity generation and dissipation.

In the limit that the protoplanets' mass is small and its Roche radius is smaller than the disk thickness, the tidal potential barrier is shallow and modest surface density variation would provide adequate pressure gradient to overcome the tidal potential barrier while preserving vortensity along the stream line. However, in cold disks with thicknesses less than the protoplanets' Roche radius, a very large surface density gradient is required to create the critical pressure gradient force needed to overcome the tidal barrier along the stream line. Consequently, mass supply from the feeding zone into the vicinity of the protoplanet is quenched. In a pressure-free gas disk, the horseshoe stream lines of cold gas cannot join onto those in the disk around the protoplanet. Only particles with a limited range of initial orbital parameters that do not follow streamlines can directly strike the protoplanet.

Based on this consideration, we suggest that the asymptotic mass of the protoplanet is determined by the condition $R_R \sim H$. This condition is similar to the thermal criterion for gap formation (Lin & Papaloizou 1986). When the planet approaches this value, the




Fig. 17.— The steady-state gas accretion rate as a function of M_p for models series 4 and 5 are represented by solid and dotted lines respectively. Both simulations show a significant decrease in $\dot{M}$, but it occurs at much lower mass for the low temperature nebula.




Fig. 18.— The planet’s mass after 20 Myr of gas accretion, starting with $M_p = 30M_\oplus$ initially. The gas depletion time scale is set to be $\tau_{\text{dep}} = 3$ Myr. The solid line designates the model series with $T_{\text{neb}} = 50K$, while the dotted line is for the model series with $T_{\text{neb}} = 100K$. As suggested from Figure 17, and seen in Figure 2, the asymptotic mass of the lower temperature model series is significantly lower than in the high T_{neb} model series.

mass supply into the gap is much reduced and the dynamics of the gas is slightly altered (Bryden et al. 1999). Although these consequence do not directly terminate protoplanets’ growth, they contribute to the reduction of the accretion rate onto the protoplanet.

The gaseous material which enters into the protoplanet’s Roche lobe forms a circumplanetary disk. Angular momentum is transferred from the circumplanetary disk by both tidally induced spiral shocks and turbulent viscosity. Initially, the flux through the circumplanetary disk is high and the disk environment is prohibitive for the formation of regular satellite systems (Lunine & Stevenson 1982; Canup & Ward 2002; Mosqueira & Estrada 2003; Canup & Ward 2006). However, our results indicate that during the final stage of the protoplanet’s growth, the characteristic timescale for protoplanetary accretion increases beyond the circumstellar disk depletion timescale. It is at this stage that physical environment in the circumplanetary disks become less hostile to satellite formation (Canup & Ward 2006). The studies of satellite formation process is beyond the scope of the present paper and they will be presented elsewhere.

In the final analysis, we are interested in the origin of the present-day mass distribution of extrasolar planets which, is observed to be $dN/dM \propto M_p^{-1}$ to $M_p^{-1.4}$ in the range of a fraction to several M_J (Marcy et al. 2005). This distribution also has a cut-off below $M_p \sim 10 - 20M_J$. Since the host stars of most known extrasolar planets have masses $M_* \sim M_\odot$, the corresponding $q = M_p/M_*$ has a similar distribution $dN/dq \sim q^{-1}$ to $q^{-1.4}$ in the range $\sim 10^{-3}$.

Scaling the formula derived in §2.3 to the values used in this simulation and assuming a dissipation rate of $\tau_{dep} = 3 - 5\text{Myr}$ and $\beta = 1$, M_p is comparable to the mass of Saturn for an aspect ratio of 0.04 and that of Jupiter for an aspect ratio of 0.07. The most favorable location for the formation of gas giant planets is near the snow line a_{ice} (Ida & Lin 2004). The magnitude of a_{ice} is determined by both the gas accretion rate within the disk and the intensity of stellar irradiation and during the classical T-Tauri phase, and a_{ice} can evolve over nearly an order of magnitude in scale (Garaud & Lin 2006). The disk aspect ratio at the snow line ($H(a_{ice})/a_{ice}$) can also change by a factor of two. Depending on the epoch of gas-giant planet formation, the variation in ($H(a_{ice})/a_{ice}$) can lead to an order of magnitude spread in the asymptotic mass of the planets. Thus, the observed planetary mass distribution may be used as to infer the ratio of the planet formation and disk depletion timescales. Detailed simulations of such models will be presented elsewhere.

Target lists for planet searches using the radial velocity technique have recently been expanded to include M-dwarf stars. However, the detection frequency for Jupiter mass planets around these low-mass stars appears to be much lower than that around G dwarfs. Until recently, the only known member of this population with known Jupiter-mass planets

was GJ 876. The lack of massive gas-giants is also conspicuous in the microlensing survey for extrasolar planets (Beaulieu et al. 2006). The difficulty for gas-giant planet formation may be due to the inadequate amount of heavy elements for assembling sufficiently massive protoplanetary cores (Ida & Lin 2004) or inability to overcome type I migration and retain sufficiently mass cores to initiate the process of rapid gas accretion (Ida & Lin 2006). In this paper, we show that even in the low-probability event that proto gas-giant planets may form, their asymptotic mass may severely limited by the tidal barrier in accordance with Equation (33). If the surface densities of both gas and dust increases with $M_\star^2$ as suggested by observationally inferred $\dot{M}$'s from the disks onto their central stars, the onset of dynamical gas accretion would occur at much later epoch with a relatively small total gas content around M dwarfs than around solar type stars. At this stage, the value of $H(a_{\text{ice}})/a$ is also an slowly increasing function of $M_\star$. All of these effects combine sensitively to quench the gas accretion rate and severely limit the asymptotic mass of the planets well below that of Jupiter.

The co-existence of multiple gas-giant planets around the same host star suggests that their asymptotic mass is limited by local tidal truncation (through gap formation) rather than the global depletion of the disk. The asymptotic mass distribution within such systems is determined by both the planets' formation location and epoch. In a steady-state disk, the value of the aspect ratio H/a determined from the disk mid-plane structure generally increases with a (Garaud & Lin 2006). In a viscous evolving disk, Σ in the outer regions of the disk decreases with a much more rapidly than in a steady disk. In general, H/a attains a maximum at a radial location a_{max} which is an increasing function of the the accretion rate through the disk. During the depletion of the disk, a_{max} decreases. Outside a_{max} , the disk is shielded from the stellar irradiation and the density scale decline rapidly with radius. This process may determine the mass distribution within multiple gas giant planet systems. It may also limit the domain of gas-giant formation.

We thank Dr. P. Bodenheimer, S. Ida, and Hui Li for useful conversation and Dr. Geoff Bryden for making available a 2D numerical hydrodynamic scheme. This work was supported in part by NASA (NAGS5-11779, NNG06-GF45G, NNG04G-191G, NNG05-G142G), JPL (1270927), NSF (AST-0507424), IGPP, and the California Space Institute.

REFERENCES

- Aarseth, S. J., Lin, D. N. C., & Palmer, P. L. 1993, *ApJ*, 403, 351
- Adachi, I., Hayashi, C., & Nakazawa, K. 1976, *Progress of Theoretical Physics*, 56, 1756

- Armitage, P. J. 1998, *ApJ*, 501, L189+
- Artymowicz, P. & Lubow, S. H. 1994, *ApJ*, 421, 651
- Balmforth, N. J. & Korycansky, D. G. 2001, *MNRAS*, 326, 833
- Bate, M. R., Lubow, S. H., Ogilvie, G. I., & Miller, K. A. 2003, *MNRAS*, 341, 213
- Beaulieu, J.-P., Bennett, D. P., Fouqué, P., et al. 2006, *Nature*, 439, 437
- Bodenheimer, P. & Pollack, J. B. 1986, *Icarus*, 67, 391
- Bryden, G., Chen, X., Lin, D. N. C., Nelson, R. P., & Papaloizou, J. C. B. 1999, *ApJ*, 514, 344
- Bryden, G., Lin, D. N. C., & Ida, S. 2000, *ApJ*, 544, 481
- Canup, R. M. & Ward, W. R. 2002, *AJ*, 124, 3404
- . 2006, *Nature*, 441, 834
- D’Angelo, G., Henning, T., & Kley, W. 2002, *A&A*, 385, 647
- Frank, J., King, A., & Raine, D. 1992, in *Accretion power in astrophysics*
- Garaud, P. & Lin, D. N. C. 2006, *ArXiv Astrophysics e-prints*
- Goldreich, P. & Lynden-Bell, D. 1965, *MNRAS*, 130, 125
- Goldreich, P. & Tremaine, S. 1979, *ApJ*, 233, 857
- . 1980, *ApJ*, 241, 425
- Greenzweig, Y. & Lissauer, J. J. 1990, *Icarus*, 87, 40
- Guillot, T. 2005, *Annual Review of Earth and Planetary Sciences*, 33, 493
- Haisch, K. E., Lada, E. A., & Lada, C. J. 2001, *ApJ*, 553, L153
- Hartmann, L. 2005, in *ASP Conf. Ser. 337: The Nature and Evolution of Disks Around Hot Stars*, ed. R. Ignace & K. G. Gayley, 3–+
- Hayashi, C., Nakazawa, K., & Nakagawa, Y. 1985, in *Protostars and Planets II*, ed. D. C. Black & M. S. Matthews, 1100–1153
- Ida, S. & Lin, D. N. C. 2004, *ApJ*, 604, 388

- . 2006, in preperation
- Klahr, H. & Kley, W. 2006, *A&A*, 445, 747
- Kley, W. 1989, *A&A*, 208, 98
- . 1999, *MNRAS*, 303, 696
- Kley, W., D’Angelo, G., & Henning, T. 2001, *ApJ*, 547, 457
- Kley, W. & Hensler, G. 1987, *A&A*, 172, 124
- Kokubo, E. & Ida, S. 1998, *Icarus*, 131, 171
- Korycansky, D. G. & Papaloizou, J. C. B. 1996, *ApJS*, 105, 181
- Laughlin, G., Steinacker, A., & Adams, F. C. 2004, *ApJ*, 608, 489
- Li, S. & Lin, D. 2006, in preparation
- Lin, D. N. C., Bodenheimer, P., & Richardson, D. C. 1996, *Nature*, 380, 606
- Lin, D. N. C. & Papaloizou, J. 1979, *MNRAS*, 188, 191
- Lin, D. N. C. & Papaloizou, J. 1985, in *Protostars and Planets II*, ed. D. C. Black & M. S. Matthews, 981–1072
- . 1986, *ApJ*, 309, 846
- Lin, D. N. C. & Papaloizou, J. C. B. 1993, in *Protostars and Planets III*, ed. E. H. Levy & J. I. Lunine, 749–835
- Lissauer, J. J. 1987, *Icarus*, 69, 249
- Lubow, S. H., Seibert, M., & Artymowicz, P. 1999, *ApJ*, 526, 1001
- Lunine, J. I. & Stevenson, D. J. 1982, *Icarus*, 52, 14
- Marcy, G., Butler, R. P., Fischer, D., et al. 2005, *Progress of Theoretical Physics Supplement*, 158, 24
- Miki, S. 1982, *Progress of Theoretical Physics*, 67, 1053
- Mosqueira, I. & Estrada, P. R. 2003, *Icarus*, 163, 198
- Nelson, R. P. & Papaloizou, J. C. B. 2003, *MNRAS*, 339, 993

- Palmer, P. L., Lin, D. N. C., & Aarseth, S. J. 1993, *ApJ*, 403, 336
- Papaloizou, J. C. B. & Lin, D. N. C. 1989, *ApJ*, 344, 645
- Papaloizou, J. C. B. & Terquem, C. 2006, *Reports of Progress in Physics*, 69, 119
- Pollack, J. B., Hubickyj, O., Bodenheimer, P., et al. 1996, *Icarus*, 124, 62
- Safronov, V. S. & Zvjagina, E. V. 1969, *Icarus*, 10, 109
- Silverstone, M. D., Meyer, M. R., Mamajek, E. E., et al. 2006, *ApJ*, 639, 1138
- Tanigawa, T. & Watanabe, S.-i. 2002, *ApJ*, 580, 506
- van Leer, B. 1977, *Journal of Computational Physics*, 23, 276
- Ward, W. R. 1998, *Bulletin of the American Astronomical Society*, 30, 1057